

Semi-inclusive spin asymmetries in polarised deep inelastic electron scattering

E.E.W. Bruins for the HERMES Collaboration

*Laboratory for Nuclear Science, Massachusetts Institute of Technology,
77 Massachusetts Avenue, Cambridge, MA 02139-4307
Temporary Address: DESY, Notkestrasse 85, 22607 Hamburg, Germany*

ABSTRACT

In 1995, the HERMES experiment at DESY (Hamburg, Germany) has taken its first data, using the 27.5 GeV polarised HERA positron beam impinging on a longitudinally polarised internal ^3He target. Preliminary results from the semi-inclusive physics program are presented.

Polarised deep-inelastic lepton-nucleon scattering has provided information about the spin structure of the nucleon, which can be expressed in terms of spin distribution functions of partons in the infinite momentum frame.

Up to now, most information has come from inclusive measurements, detecting the scattered lepton only. It is expected that the investigation of the hadronic final state in the scattering process will allow a deeper insight into the relevant processes than the inclusive measurement alone. In HERMES [1], the scattered positron and the hadronic final state are measured in a wide kinematic range and with a good angular and momentum resolution. Positrons have been clearly identified and separated from the hadrons using preshower, calorimeter and Čerenkov counter information. Semi-inclusive pions were identified and separated from heavier hadrons by means of the Čerenkov counter (above 5.6 GeV/c) and time-of-flight techniques (below 2 GeV/c). Semi-inclusive final state hadrons which decay in the detector volume have been identified by the reconstruction of the invariant mass of




Figure 1: Spin asymmetry of semi-inclusive, positively charged hadrons, for the polarised H, D (SMC) and ^3He (HERMES, preliminary)

the decaying system. A full description of the experiment, the detector set up and the target can be found in [2].

The analysis of the 1995 semi-inclusive data is in progress. We focus on both the polarised and unpolarised physics potential of HERMES, and expect to deliver the first complete quark spin decomposition on the polarised neutron. In Figure 1 the ^3He spin asymmetry is shown for positively charged semi-inclusive hadrons $A_1^{h^+}(^3\text{He})$ for $z = p/\nu > 0.2$, where p is the momentum of the hadron and ν the energy transfer in the reaction. The data indicate a small negative asymmetry, in agreement with the naive expectation based on charge/isospin conjugation and existing SMC data for polarised H and D , measured on polarised butanol and deuterated butanol targets [3]. Monte Carlo studies with PEPSI [4], based on the LUND fragmentation model, have shown that, due to the forward acceptance of the HERMES detector, an even lower z -cut does not significantly change the underlying physics in our data. In Figure 2 we show the spin asymmetry for semi-inclusive neutral pions, identified by the reconstruction of the invariant mass of the two decay gammas.

Due to the relatively high Čerenkov pion threshold (5.6 GeV/c) during the 1995 run, the spin asymmetry for *identified* charged pions has large statistical uncertainties. The 1996 data on the proton will profit from a lower threshold, and lead to a significant improvement in our semi-inclusive pion data. Using linear combinations of the various semi-inclusive spin asymmetries and (model dependent) Monte Carlo predictions, we thus expect to extract quark spin distributions as a function of x_{Bj} from the ^3He and proton data, separately. Finally, the combination of the 1995 and 1996 data will allow an extraction of the valence quark spin distributions Δu_v and Δd_v as a function of x_{Bj} .




Figure 2: Spin asymmetry of semi-inclusive, neutral pions for polarised ^3He (HERMES, preliminary). The curve shows a parton model prediction using the Schäfer parametrisation [5].

Preliminary results for unpolarised ρ production have been extracted. These indicate the superior quality of the data which HERMES is expected to deliver in the forthcoming years. Figure 3 shows the decay angle distribution of elastic, diffrac-

tive ρ mesons, without subtraction of the (small) non-exclusive and combinatorial background. Figure 4 shows the nuclear transparency of ${}^3\text{He}$ (normalized to $H + D$ data from less than 24 hours dedicated data taking) to the incoherently produced sub-sample of the above mentioned ρ mesons. Although the statistics is low and the systematics not yet fully understood, combined with our 1996 proton data, we expect to measure the nuclear transparency of ${}^3\text{He}$ as a function of momentum and energy transfer in the reaction. Targets with $A > 3$ are being considered at the moment.

- [1] K. Coulter *et al.*, DESY-PRC 90/01 (1990); M. Düren, DESY-HERMES 95-02 (1995)
- [2] A. Simon, this symposium
- [3] B. Adeva *et al.*, SMC Collaboration, Phys. Rev. **B369** (1996) 93
- [4] G. Ingelmann, LEPTO – CERN programming pool W5045 (1986); M. Veltri, H. Ihssen, PEPSI – HERMES Collaboration (1995)
- [5] A. Schäfer *et al.*, hep-ph/9505306




Figure 3: (a) The decay angle distribution of elastic diffractive rho mesons from ${}^3\text{He}$ (No background subtraction); (b) Nuclear transparency of ${}^3\text{He}$ to incoherently produced elastic, diffractive rho mesons, as a function of Q^2 . We allow for a preliminary 20 % overall normalisation uncertainty (HERMES, preliminary)